

## **MEETING ABSTRACT**

**Open Access** 

## The pathological significance of selected *MLH1* missense mutations based on the theoretical *in silico* predictions

Mariola Dubanowicz

From Annual Conference on Hereditary Cancers 2013 Szczecin, Poland. 26-27 September 2013

Germline mutations in the human DNA mismatch repair (MMR) genes MSH2 and MLH1 are associated with the inherited cancer disorder Lynch syndrome (LS). Up to 30% of MLH1 variants found are missense mutations. The functional consequences in regard to pathogenicity of many of these variants are unclear. Missense mutations affect protein structure or function, but may also cause aberrant splicing. In this study, in silico prediction tools (SIFT, PolyPhen-2 and ESEfinder 3.0), literature and an online MMR mutation database review were used to evaluate the clinical significance of 26 MLH1 missense mutations identified in patients from Polish families, suspected of Lynch syndrome. Six MLH1 VUS were classified as pathogenic, nine MLH1 missense mutations are classified as neutral. 11 variants require additional research. The results show that in silico prediction tools can be utilized in an appropriate and efficient manner to determine the pathogenicity of MMR gene variations.

Published: 9 September 2015

doi:10.1186/1897-4287-13-S1-A9

Cite this article as: Dubanowicz: The pathological significance of selected *MLH1* missense mutations based on the theoretical *in silico* predictions. *Hereditary Cancer in Clinical Practice* 2015 13(Suppl 1):A9.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



Pomeranian Medical University, Faculty of Medical Biotechnology, Szczecin, Poland

